ALLERGIES AND ALLERGIC REACTIONS

Allergies occur when the **immune system** identifies and **reacts** to a **foreign substance** such as food, **environmental allergens** (*pollen*, *pet dander*, *insect venom*) and **medications**.

Listed below are different types of allergies and the symptoms they can cause.



(aka: hayfever, **seasonal allergies**, **pollen** allergy, **pet** allergy, etc.)

- Sneezing
- Itching of the nose, eyes or roof of the mouth
- Runny, stuffy nose
- Watery, red or swollen eyes (conjunctivitis)

Food Allergy

(common causes are **peanuts** or tree nuts, **shellfish**, milk, eggs, soy)

- **Tingling** in the mouth
- Swelling of the lips, tongue, face or throat
- Hives
- Anaphylaxis

Atopic dermatitis

An allergic skin condition also called **eczema**, causes skin to:

- Itch
- Redden
- Flake or peel

Check out the latest updates and details from the Mayo Clinic:



Drug Allergy

(commonly caused by antibiotics like **penicillins** or **sulfa** drugs, but could be any medication)

- Hives, rash
- Itchy skin
- Facial swelling
- Wheezing
- Anaphylaxis

Insect Sting Allergy

(commonly caused by bees)

- A large area of swelling (edema) at the sting site
- Itching or hives all over the body
- Cough, **chest tightness**, **wheezing** or shortness of breath
- Anaphylaxis

Anaphylaxis

- Facial swelling
- Severe shortness of breath, wheezing, difficulty breathing
- Skin rash, itching, or pale skin
- A drop in blood pressure, causing:
- Lightheadedness, dizziness
- Fainting or loss of consciousness
- A rapid, weak pulse
- Nausea and vomiting





Treatment of minor symptoms

- **Antihistamines** (like Claritin, Zyrtec, Allegra, or Benadryl)
- Decongestants (like Afrin or Sudafed)
- Ice and topical creams that contain corticosteroids (like hydrocortisone ointment)

If someone you know is experiencing **anaphylaxis**:

- Call 911 immediately.
- See if they have an (EpiPen), epinephrine (adrenaline) autoinjector, help them if needed.
- Keep the person **calm**.
- Lie them on their **back**.
- Raise their feet about 12 inches and cover them with a blanket.
- Turn them on their side if they are vomiting or bleeding.
- Make sure their clothing is loose so they can breathe.

THE SOONER THE PERSON GETS THEIR EPINEPHRINE, THE BETTER.

Saint Liam Hall, Notre Dame (574) 631-7497 - uhs.nd.edu